

## Quantum Frog (quantum)

Beatrice is a researcher at the Italian Tandem Accelerator Collision Particles Centre and she is on the verge of a major breakthrough. She recently discovered the new Frog particle and she is now studying its behaviour in a closed system.

The system can be described by  $N$  possible energy levels  $a_i$ . From her observations so far, Beatrice has gathered the following information about the Frog particle:

- It has quantum properties: it is located at two different energy levels at the same time, i.e. its state can be represented by a pair  $(i, j)$ .
- It has frog properties: it can jump from one energy level to another, but it cannot jump from both energy levels at the same time, i.e. in one jump from  $(i, j)$  it can land on  $(i, k)$  or  $(h, j)$  but not on  $(h, k)$ .

However, not all jumps are possible: Beatrice knows that the entropy of the system must be strictly increasing after each jump. For an isolated Frog particle in state  $(i, j)$ , she determined the entropy to be equal to the Frog constant  $F$  multiplied by the absolute difference  $|a_i - a_j|$  between the energy levels of the state.

Beatrice performed one observation on the system and she identified the Frog particle in state  $(x, y)$ . After the observation, the system naturally evolved and the Frog particle performed an unknown sequence of (possibly zero) entropy-increasing jumps. To help Beatrice complete her research, you should compute the number of possible distinct states where the Frog particle may now be located (the state  $(i, j)$  is considered equivalent to the state  $(j, i)$ ).



Figure 1: What society thinks Beatrice is doing.

### Input

The first line contains three integers  $N$ ,  $x$ ,  $y$ , respectively, the number of energy levels of the system and the state where the Frog particle was located during the observation.

The second line contains  $N$  integers  $a_0, a_1, \dots, a_{N-1}$ , the values of the energy levels.

### Output

You should output a single line containing a single integer: the number of possible distinct states that the Frog particle may have reached after the observation.

## Constraints

- $2 \leq N \leq 5 \cdot 10^5$ .
- $0 \leq x, y < N$ .
- $x \neq y$ .
- $0 \leq a_i \leq 10^{18}$  for every  $i$ .
- $a_i \neq a_j$  for every  $i \neq j$ .

## Examples

input	output
4 1 2 40 20 30 10	4
5 0 1 10 20 30 40 50	6

## Explanation

In the **first sample case**, the Frog particle can either be in states  $(0, 1)$ ,  $(1, 2)$ ,  $(2, 3)$  or  $(0, 3)$ .

In the **second sample case**, the Frog particle can either be in states  $(0, 1)$ ,  $(0, 2)$ ,  $(0, 3)$ ,  $(0, 4)$ ,  $(1, 3)$  or  $(1, 4)$ . Note that although the state  $(2, 4)$  has an entropy strictly greater than the initial one, it cannot be reached with any sequence of jumps.